

# Co-Actyl H



A biostimulant organic complex with 3 main actions for better development

## 1. Activated Humic and Fulvic Acids - Crop nutrition improvement by better availability of soil nutrients.

H.F.A. selected are treated under a partial **solubilization process** making them **120 times** more soluble than H.F.A. on the market.

Therefore activated H.F.A. are immediately dispersed in the soil solution to form stable and soluble humo-complex with nutrients which is difficult for the plant under normal circumstances.

## 2. Zeatin - superb growth stimulation, increasing the root functions and root mass.




### 3. Phytoalexin Complex

The formulation of Three active components determined by the Department of Vegetal Physiology of the University of Navarre (Spain) and the Roullier Research Center. The actives stimulate the production of ALEXINES within the plant which in turn improves the resistance to pathogens

#### Phytoalexin Complex

Co-Actyl-H stimulates the message molecules activating more Phytoalexine protein production



#### Special for HLB Greening Applications:

**Dosage:** Apply for each application via herbicide boom to soil (weeds not present) 15 lbs. Per treated acre. It should be applied 15 to 40 days prior to a major fertilizer application. It should be used 3 times per year. Preferably applied in late February, Early March then late May, early June, and last application in September. Application must be followed immediately with 1/2 inch irrigation. Microjet irrigation can be running while spray is being applied. Do not apply with acidic fertilizers or calcium nitrate. Application by itself is recommended.

USE 60 mesh strainer on Tank strainer and 50 mesh strainers on nozzle tips.

Not compatible with acid products or products containing calcium, magnesium and/or soluble salt of microelements (Fe, Cu, Mn, Zn and B).

#### Special for foliar or drip irrigation applications (watersoluble)

**Dosage:** Apply 40 to 60 lbs. Per Acre. Can be applied 2 to 3 times for growing season. Foliar application must be followed immediately with 1/2 inch irrigation. Can be used with good results in Dry Jet applicators. Do not apply with acidic fertilizers or calcium nitrate. Application by itself is recommended. USE 60 mesh strainer on Tank strainer and 50 mesh strainers on nozzle tips.

#### Effectiveness:

- Stimulates fast growth of feeder roots
- Improves Soil to Plant relationship
- Increases Plant's Natural Defenses
- Nutrition from soil is evident with growth.